


Recall:

For any angle of interest (θ), there are three (3) primary trigonometric ratios.

$$\text{sine of } \theta = \frac{\text{opposite}}{\text{hypotenuse}}$$

$$\text{cosine of } \theta = \frac{\text{adjacent}}{\text{hypotenuse}}$$

$$\text{tangent of } \theta = \frac{\text{opposite}}{\text{adjacent}}$$





S o h C a h T o a

Apr 25-9:54 PM

Radian Angles on the Cartesian Plane

The Special Triangles can be used to identify exact values for trigonometric ratios of special angles.




$$\sin \frac{\pi}{3} = \frac{\sqrt{3}}{2}$$

Oct 23-9:32 PM

Radian Angles on the Cartesian Plane

An angle is in standard position if the vertex is at the origin and the **initial arm** is along the positive x-axis.

This angle can be described in terms of the point (x, y) at the end of the **terminal arm**,




where: $r^2 = x^2 + y^2$

$$\sin \theta = \frac{y}{r} \quad \cos \theta = \frac{x}{r} \quad \tan \theta = \frac{y}{x}$$

$$\csc \theta = \frac{r}{y} \quad \sec \theta = \frac{r}{x} \quad \cot \theta = \frac{x}{y}$$

Apr 25-10:21 PM

The related acute angle (RAA) is the positive, acute angle between the nearest x-axis and the terminal arm.



$$0^\circ < \text{RAA} < 90^\circ$$

$$0 < \text{RAA} < \frac{\pi}{2}$$

$$\begin{aligned} \pi - \text{RAA} &= \frac{5\pi}{6} \\ \text{or} \quad \pi - \frac{5\pi}{6} &= \text{RAA} \\ \text{RAA} &= \frac{\pi}{1} - \frac{5\pi}{6} \\ &= \frac{6\pi - 5\pi}{6} \end{aligned}$$

$$\boxed{\text{RAA} = \frac{\pi}{6}}$$

Oct 23-8:49 AM

The CAST rule allows us to quickly determine the sign of each trig ratio for any quadrant.

| | | | |
|----------------------|---------------------|--|--|
| Q2 | | Q1 | |
| $-x$ $+y$ $+r$ | sine positive | all positive $+x$ $+y$ $+r$ | $+sin$ $-cos$ $-tan$ S |
| $-x$ $-y$ $+r$ | tangent positive | cosine positive $+x$ $-y$ $+r$ | $+sin$ $+cos$ $+tan$ A |
| Q3 | | Q4 | |
| | | $-sin$ $-cos$ $+tan$ T | $-sin$ $+cos$ $-tan$ C |

Use the CAST rule, along with the Related Acute Angle (RAA) to solve for the angle.

May 3-9:19 AM

Ex.1 Evaluate using Cartesian definitions & special triangles.

(a) $\sin\left(\frac{\pi}{2}\right)$

(b) $\cos\left(\frac{5\pi}{4}\right)$

(a) $\sin \frac{\pi}{2} \left[\sin \theta = \frac{y}{r} \right]$

$= \frac{1}{1}$ \leftarrow y-coord
 \leftarrow r-value

$= 1$




(b) $\cos\left(\frac{5\pi}{4}\right)$

$= -\cos\left(\frac{\pi}{4}\right)$

\uparrow CAST \uparrow RAA

$= -\left(\frac{1}{\sqrt{2}}\right)$

$= \frac{-1}{\sqrt{2}}$



$\frac{s}{T} \mid \frac{A}{C}$

Oct 23-11:07 PM

Ex.2 Solve $\tan \theta = \frac{-7}{24}$ for $0 \leq \theta < 2\pi$

① CAST $\begin{array}{c|c} \text{S} & \text{A} \\ \hline \text{T} & \text{C} \end{array}$


* radian mode
on
calculator.

Q2 or Q4

② solve for RAA: $\tan(\text{RAA}) = \frac{7}{24}$


$$\begin{aligned} \text{RAA} &= \tan^{-1}\left(\frac{7}{24}\right) \\ &\approx 0.2838 \text{ rad} \end{aligned}$$

③ Q2



$$\begin{aligned} \theta + \text{RAA} &= \pi \\ \theta &= \pi - \text{RAA} \\ \theta &\approx 2.8578 \end{aligned}$$

Q4



$$\begin{aligned} \theta + \text{RAA} &= 2\pi \\ \theta &= 2\pi - \text{RAA} \\ \theta &\approx 5.9994 \end{aligned}$$

check: $\tan \theta = \frac{-7}{24}$

$$\begin{aligned} \text{LS} &\approx \tan(2.8578) \\ &\approx -0.2917 \end{aligned}$$

$$\begin{aligned} \text{RS} &= \frac{-7}{24} \\ &\approx -0.2917 \end{aligned}$$

Oct 23-11:11 PM

Assigned Work:

p.330 # 1-4, 5ace, 6ace, 7ace, 8ace, 9, 11, 13

Oct 23-10:59 PM