

Factoring Strategies

Nov 1/2011

Consider the factoring methods we have explored so far:

- \* 1. Common Factors  $3x^2 + 6x + 9$
- \* 2. Factoring by Grouping  $ac + ad + bc + bd$
- 3. Simple Trinomials  $1x^2 + bx + c$
- \* 4. Complex Trinomials  $ax^2 + bx + c, a \neq 1$  SPI
- 5. Perfect Squares  $a^2 + 2ab + b^2$
- 6. Difference of Squares  $a^2 - b^2$

It is often sufficient to use only one method, but there are times when they must be combined. This occurs most often when **common factors** are involved.

Mar 29-11:14 AM

Always check for common factors before you start **and** after you think you are done.

When you are asked to "fully factor" or "factor completely", all common factors must also be accounted for.

Ex.1 Remove Common Factors First and Last

$$4x^2 - 20x + 24$$

$$\begin{array}{l|l}
 = 4(x^2 - 5x + 6) & = 4x^2 - 8x - 12x + 24 \quad \begin{array}{l} S: -20 \\ P: 96 \\ I: -8, -12 \end{array} \\
 = 4(x-2)(x-3) & = 4x(x-2) - 12(x-2) \\
 & = (x-2)(4x-12) \\
 & = (x-2)(4)(x-3) \\
 & = 4(x-2)(x-3)
 \end{array}$$

Oct 31-10:46 PM

Ex.2 Determine the <sup>best</sup> strategies required to factor:

(a)  $x^2 + 8x + 15$

simple

(d)  $4x^2 - 8x + 4$

common  
perfect? / simple

(b)  $6x^2 + 19x + 8$

complex

(e)  $9x^2 + 48x + 64$

perfect sq.

(c)  $40x^2 - 250$

common factor  
diff. of squares

(f)  $-5x^2 + 60x - 180$

common factor -5  
simple

Mar 29-11:16 AM

Assigned Work:

p. 236 # 1, (6-8)ace, 9, 10, 12, 14ac, 17\*

Mar 26-9:06 AM

10(e)

$$2(a+b)^2 + 5(a+b) + 3$$

$\underbrace{\hspace{2cm}}_x$        $\underbrace{\hspace{2cm}}_x$

$$= 2x^2 + 5x + 3$$

$$= 2x^2 + 3x + 2x + 3$$

$$= x(2x+3) + 1(2x+3)$$

$$= (2x+3)(x+1)$$

$$= (2(a+b)+3)(a+b+1)$$

$$= (2a+2b+3)(a+b+1)$$

S 5

P 6

I 3, 2

Nov 2-10:29 AM

6(e)  $12x^2 + 4x - 21$

$$= 12x^2 - 14x + 18x - 21$$

$$= 2x(6x-7) + 3(6x-7)$$

$$= (6x-7)(2x+3)$$

S 4

P -252

I

-1 252

-2 126

-3 84

-4 63

⋮

-14 18 ✓

FOIL

$6x - 7$		
$2x + 3$		

Nov 2-10:33 AM

$$\begin{aligned}
 7(c) \quad & 18x^2 + 60x + 50 \\
 & = 2(9x^2 + 30x + 25) \\
 & = 2(3x + 5)^2
 \end{aligned}$$

$$S \ 60$$

$$P \ 900$$

$$I \ 30, 30$$

check

$$\begin{aligned}
 & 2(3x)(+5) \\
 & = +30x \quad \checkmark
 \end{aligned}$$

Nov 2-10:39 AM

$$12. \ A = \pi x^2 + 10\pi x + 25\pi$$

$$(a) \ r = \underline{\quad} ?$$

$$A = \pi r^2$$

$$\frac{\pi r^2}{\pi} = \frac{\pi x^2}{\pi} + \frac{10\pi x}{\pi} + \frac{25\pi}{\pi}$$

$$r^2 = x^2 + 10x + 25$$

$$r^2 = (x+5)^2 \quad (x+5)(x+5)$$

$$r = \pm(x+5), \text{ but } r > 0$$

$$r = x+5$$

$$\begin{aligned}
 (b) \quad x=10, \quad r &= 10+5 \\
 &= 15
 \end{aligned}$$

Nov 2-10:43 AM