

Mathematical Operations in Java

Basic Mathematical Operations

Many mathematical expressions in Java look identical to those that you would write on paper. There are five basic arithmetic operators in Java, as shown below.

| Java Operator | Purpose | Example |
|---------------|--|-------------------------|
| + | Addition | $5 + 3.4 = 8.4$ |
| - | Subtraction | $203.9 - 9.12 = 194.78$ |
| * | Multiplication | $3 * 4.7 = 14.1$ |
| / | Division | $17.889 / 6.7 = 2.67$ |
| % | Modulo Division (calculates the remainder) | $35 \% 9 = 8$ |

In general, operations on values of the same type will produce an answer that is also of the same type. It is particularly important to keep this in mind when performing an *integer division* (that is, a division between two integer value, or variables containing integers). Consider the following examples:

- (a) $15 / 3$ gives 5
- (b) $13 / 4$ gives 3 (rather than 3.25)
- (c) $9 / 5$ gives 1 (rather than 1.8... notice that the answer is *truncated*, not rounded)

Whenever integer and floating point values are mixed in an expression, the result will be a floating point value. Notice the subtle difference this creates when applied to our previous example.

- (a) $15 / 3.0$ gives 5.0 (a floating point answer)
- (b) $13.0 / 4$ gives 3.25
- (c) $9 / 5.0$ gives 1.8

Mathematical Operations in Java

Modulo Division

It is worth giving an extra moment to consider and understand the *modulo operator*. This operator is used to find the *remainder* from an *integer division*, which is actually the first type of division taught to children in elementary school.

Consider the example of $35 \% 9 = 8$. To begin, we ask, “How many times does 9 divide evenly into 35?” Simply following the nine-times-table yields

$$9 \times 1 = 9 \quad 9 \times 2 = 18 \quad 9 \times 3 = 27 \quad 9 \times 4 = 36$$

The answer is obviously 3 times. Now, what is the remainder? Subtract 27 from 35 for a remainder of 8.

This mathematical concept is actually quite useful and frequently used in programming, which is why many programming languages, including Java, provide it in a compact and convenient form.

Division by Zero

Both regular division and modulo division pose the risk of accidentally performing an illegal mathematical operation, which is *division by zero*. In mathematics, division by zero is *undefined*, and this must apply to computer programs as well.

Since most calculations in programs use *variables*, it may not be obvious when or if this is going to happen. It is always a good idea to carefully consider the possible values of any variables when you perform a division operation.

If a division by zero occurs while performing an integer operation, the Java program will exit with an error. This is known as **throwing an exception**, and in this case, the exception is an **ArithmeticException**. Basically, the program is informing the user that it has encountered a situation that it doesn't know how to handle. Later we will learn about handling various types of exceptions, but for now, we will simply try to avoid this situation.

When performing floating point operations, the division by zero is handled by Java using one of three special floating point expressions, as summarized in the table below.

| Floating Point Operation | Result | Can be used in further calculations? | Will print as the string... |
|--------------------------------|-------------------|--------------------------------------|-----------------------------|
| positive value divided by zero | positive infinity | yes | Infinity |
| negative value divided by zero | negative infinity | yes | -Infinity |
| zero divided by zero | not a number | no | NaN |

Mathematical Operations in Java

Mathematical Methods

To allow for mathematical operations beyond basic arithmetic, Java provides additional functionality using a large group of *methods* contained in the *Math* class. To use any of these methods, we must identify them as being part of the *Math* class. Some of the more common and useful methods from the *Math* class are shown in the following table.

| Mathematical Operation | Java Syntax | Description |
|------------------------|-----------------------------|----------------------------|
| \sqrt{x} | <code>Math.sqrt(x)</code> | square root of x |
| x^y | <code>Math.pow(x, y)</code> | x raised to the exponent y |
| $ (x) $ | <code>Math.abs(x)</code> | absolute value of x |

Some other important mathematical methods do not correspond to traditional mathematical operations, but are very useful in programming.

| Java Syntax | Description |
|----------------------------|--|
| <code>Math.round(x)</code> | rounds a floating point value, x, to the nearest integer (up or down) |
| <code>Math.ceil(x)</code> | rounds a floating point value up to an integer value |
| <code>Math.floor(x)</code> | rounds a floating point value down to an integer value |
| <code>Math.random</code> | generates a random number between 0 and 1 ($0 \leq x < 1$) includes the value 0, but not the value 1 (0 to 0.999999...) |

Mathematical Operations in Java

Exercises

1. Evaluate each expression, and then check your answer using Java. If the answer is a floating point value (decimal), add a decimal to your answer (e.g., if the answer is 3, but a floating point, write it as 3.0). If the expression is not valid, explain why and determine how you could make the expression valid.

- (a) $2 + 7 / 2$
- (b) $5 (6 + 3.0)$
- (c) $3 * 5 / 2$
- (d) $3.0 * 5 / 2$
- (e) $2 * (3 / 4)$
- (f) $2.0 + 7 / (3 / 4)$
- (g) $3 * 4 / 5$
- (h) $35 / (5 / 4)$
- (i) $4 - [6 - (2 - 7)]$
- (j) $3 / 4 * 2.0$
- (k) $3 / (4 * 2.0)$

2. Write the following calculations in Java, paying close attention to order of operations. Use brackets, where necessary, to ensure calculations occur in the desired order. Try not to overuse brackets. Verify your Java answers with a calculator and confirm results with another member of the class.

- (a) $\frac{2+3}{2-3}$
- (b) $5+3.0 \times \frac{2}{4.0}$
- (c) $\frac{3+5}{2} + \frac{7}{5-2}$
- (d) $\frac{2.0 + \frac{3.0}{4.0}}{7.0 - 3.0}$

3. Write the following calculations in Java using the mathematical methods.

- (a) $\sqrt{25}$
- (b) $\sqrt{11}$
- (c) 3^2
- (d) 2^3