


Recall:

For any angle of interest ( $\theta$ ), there are three (3) primary trigonometric ratios.

$$\text{sine of } \theta = \frac{\text{opposite}}{\text{hypotenuse}}$$

$$\text{cosine of } \theta = \frac{\text{adjacent}}{\text{hypotenuse}}$$

$$\text{tangent of } \theta = \frac{\text{opposite}}{\text{adjacent}}$$





S o h C a h T o a

Apr 25-9:54 PM

### Radian Angles on the Cartesian Plane

Oct 24/2014


The Special Triangles can be used to identify exact values for trigonometric ratios of special angles.



Oct 23-9:32 PM

An angle is in standard position if the vertex is at the origin and the **initial arm** is along the positive x-axis.

This angle can be described in terms of the point  $(x,y)$  at the end of the **terminal arm**,



where:  $r^2 = x^2 + y^2$


$$\sin \theta = \frac{y}{r} \quad \cos \theta = \frac{x}{r} \quad \tan \theta = \frac{y}{x}$$

$$\csc \theta = \frac{r}{y} \quad \sec \theta = \frac{r}{x} \quad \cot \theta = \frac{x}{y}$$

Apr 25-10:21 PM

Ex.1 Evaluate using Cartesian definitions & special triangles.


(a)  $\sin\left(\frac{\pi}{2}\right)$       (b)  $\cos\left(\frac{5\pi}{4}\right)$



$$\sin \theta = \frac{y}{r}$$

$$\sin\left(\frac{\pi}{2}\right) = \frac{1}{1}$$

$$= 1$$



$$\cos\left(\frac{5\pi}{4}\right) = -\cos\left(\frac{\pi}{4}\right)$$

$$= -\frac{1}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}}$$

$$= -\frac{\sqrt{2}}{2}$$

Oct 23-11:07 PM

The CAST rule allows us to quickly determine the sign of each trig ratio for any quadrant.

Q2		Q1
sine positive		all positive
tangent positive		cosine positive
Q3		Q4

S		A
T		C


Use the CAST rule, along with the Related Acute Angle (RAA) to solve for the angle.

May 3-9:19 AM

Ex.2 Solve  $\tan \theta = \frac{-7}{24}$  for  $0 \leq \theta < 2\pi$

$$\tan \theta = \frac{y}{x} \quad y = -7 \quad \text{or} \quad \frac{7}{-24}$$

$$x = 24$$



$$\tan \text{RAA} = \frac{7}{24}$$

$$\text{RAA} = \tan^{-1}\left(\frac{7}{24}\right)$$

$$\doteq 0.2838$$

$$\theta_1 \doteq 2\pi - 0.2838$$

$$\doteq 5.999$$

$$\theta_2 \doteq \pi - 0.2838$$


$$= 2.8578$$

Oct 23-11:11 PM

Assigned Work:

p.330 # 1-4, 5ace, 6ace, 7ace, 8ace, 9, 11, 13

3c




$$\sin\left(\frac{7\pi}{4}\right) = -\sin\left(\frac{\pi}{4}\right) = -\frac{\sqrt{2}}{2}$$

$$\cos\left(\frac{7\pi}{4}\right) = +\cos\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2}$$


$$\tan\left(\frac{7\pi}{4}\right) = -\tan\left(\frac{\pi}{4}\right) = -1$$

Oct 23-10:59 PM



Oct 30-9:16 AM

9.



$$\sin RAA = \frac{3.4}{5}$$

$$RAA = \sin^{-1}\left(\frac{3.4}{5}\right)$$

$$\approx 0.7478$$


$$\theta = \pi - RAA$$

$$\approx 3.1416 - 0.7478$$

$$\approx 2.4$$

Oct 30-9:18 AM

11.



$60s = 2\pi$

$\frac{25}{60} \times 2\pi$


$\frac{10}{60} \times 2\pi = \frac{1}{6} \times 2\pi = \frac{\pi}{3}$

$\cos \frac{\pi}{3} = \frac{x}{9}$

$x = 9 \cos\left(\frac{\pi}{3}\right)$

$= 9\left(\frac{1}{2}\right)$

$= 4.5$



Oct 30-9:20 AM